

CHECK LIST

Preeclampsia AND H.T

- Definition = New onset of hypertension and proteinuria after 20 weeks gestation.
 - Systolic blood pressure ≥ 140 mmHg OR diastolic blood pressure ≥ 90 mmHg
 - Proteinuria of 0.3 g or greater in a 24-hour urine specimen
 - Preeclampsia before 20 weeks, think MOLAR PREGNANCY!
- Categories
 - Mild Preeclampsia
 - Severe Preeclampsia
- Eclampsia
 - Occurrence of generalized convulsion and/or coma in the setting of preeclampsia, with no other neurological condition.
- Severe Preeclampsia must have one of the following:
 - Symptoms of central nervous system dysfunction
 - Symptoms of liver capsule DISTENTION
 -
 -

-
-
-
-
-
-

Treatment of Preeclampsia

Gestational Hypertension

- Mild hypertension without proteinuria or other signs of preeclampsia.
- Develops in late pregnancy, after 20 weeks gestation.
- Resolves by 12 weeks postpartum.

Risk Factors for Hypertension in Pregnancy

Evaluation of Hypertension in Pregnancy

History

Physical

Laboratory Tests

Management of Hypertension in Pregnancy

- Medical Management
- Contraindicated antihypertensive drugs
- Proceed with Delivery
- Only cure is DELIVERY!!!